



Colour vision defect in phakic vs pseudophakic eye group

Shua Azam^{1*}, Abdul Hameed Talpur², Zeenat Bano³, Mehjabeen Jaffri⁴, Hina Haider⁵, Vineeta Kumari⁶, Sadia Bukhari⁷

¹ MPhil. Optometry, Senior Lecturer, ISRA School of Optometry, Al-Ibrahim Eye Hospital, Karachi, Pakistan

² Ophthalmologist, ISRA School of Optometry, Al-Ibrahim Eye Hospital, Karachi, Pakistan

^{3,5,6} Optometrist, Sindh Institute of Ophthalmology and Visual Sciences (SIOVS) Eye Hospital Hyderabad, Pakistan

⁴ Internee Optometrist, Al-Ibrahim Eye Hospital, Karachi, Pakistan

⁷ Professor of Ophthalmology, Al-Tibri Medical College ISRA Postgraduate Institute of Ophthalmology, Karachi, Pakistan

Abstract

Objective: To determine the color vision defect in pseudophakic and phakic group.

Material and Method: This was cross sectional with non-probability convenient sampling from June to November 2019 at Al-Ibrahim Eye Hospital, Karachi. The patient's age lied between 45 to 60 years and had a follow up after 1 month of surgery. Visual acuity taken was from 6/6 to 6/18 after refraction. All types of refractive errors after Phacoemulsification surgery (Acrylic IOL) were included. Posterior sub-capsular opacity and other types of cataract surgery were excluded. The Panel D - 15 test was used to assess color vision defect. Ethical approval was given by Research Ethical Committee (REC) of ISRA Postgraduate Institute of Ophthalmology. Statistical analysis was done by statistical package for social science (SPSS) version 20.0.

Results: A total of 160 eyes were enrolled in this study. Among 160 eyes; 70 (44%) and 90 (56%) eyes were of males and females, respectively. The eyes were categorized in two groups, 80 (50%) were phakic and 80 (50%) were pseudophakic. Among 80 (50%) pseudophakic eyes; 32 eyes had Tritanopia (40%), 10 eyes had Deuteronopia (13%), 8 eyes had Protonopia (10%), 6 eyes had combined Tritonopia and Protonopia (8%), 4 eyes had combined Deuteronopia and Tritonopia (5%), 2 eyes had Protonopia and Deuteronopia (3%) and 18 (23%) had no defect. Among 80 phakic eyes, 72 (90%) had no defect while Deuteronopia was found in 1(1.25%) eye and Tritanopia in 7 (9%) eyes.

Conclusion: Tritonopia was mostly present in Pseudophakic group while majority of Phakic group do not show color vision defect.

Keywords: pseudophakic, phakic, panel d-15, acrylic (IOL), color vision defect (CVD)

1. Introduction

Color vision deficiencies diminish the capacity to separate certain colors under specific circumstances and its testing identify the existence, type, and severity of defects, providing a basis for the evaluation of the defect's impact on personal and professional performance [1]. Color vision discrimination deteriorate with progressing age [2]. Ocular diseases such as cataract and glaucoma [3], trauma and certain medication also affect color vision [4]. Chromatic discrimination is assessed by color vision testing [5] by using different color vision tests. [6, 7, 8, 9]

In European Caucasians the prevalence of color vision deficiency is about 8% in men and about 0.4% in women and between 4% & 6.5% in men of Chinese and Japanese ethnicity, respectively [10].

Some regional prevalence studies showed diversity in prevalence such as Turkey (7.3%), Iran (4.7%), India (2.8% to 8.2%, ethnic variations) and Saudi Arabia (2.9%) [11]. While in Pakistan color vision deficiency (CVD) ranges from 0.9% [12], 2.48% [13] and 2.78% [14].

Methodology

This cross-sectional study with non-probability convenient sampling was carried out in the Male and Female OPD at

Al-Ibrahim Eye Hospital, Karachi, Pakistan. The study duration was June 2019 to November 2019. The protocol for examination for all patients who met our inclusion exclusion criteria were included. Visual acuity was recorded separately both for near and distance, with and without glasses and with pinhole. A total of 160 eyes were taken with 80 eyes pseudophakic and 80 eyes phakic. Inclusion criteria included age ranged from 45 years to 60 years old, Phacoemulsification surgery with Acrylic IOL implant, follow-up after one month & all types of refractive errors after cataract extraction and visual acuity ranges from 6/18 to 6/6. Posterior sub capsular opacity and other types of cataract surgeries and systemic diseases were excluded. The Panel D15 test was performed at 33cm distance to find the changes in color vision. Self-prepared Performa was used for collection of data. Statistical analysis was done on statistical package for social science (SPSS) version 20.0. All the categorical variables were presented as frequencies and percentages.

Results

The eyes were categorized in the two groups: Phakic group with 80 eyes and pseudophakic group with 80 eyes in a total sample of 160 eyes as shown in Table 1.

Table 1: Pseudophakic and Phakic Eyes

Groups	Frequency	Percentages %
Pseudophakic Eyes	80	50.0%
Phakic Eyes	80	50.0%
Total	160	100.0%

All included sample was examined for right and left eye separately. Among them 86 (54%) were right eyes and 74 (46%) were left eyes as shown in Table 2.

Table 2: Distribution of Eyes

Eyes	Frequency	Percentages
Right Eye	86	53.8%
Left Eye	74	46.3%
Total	160	100.0%

The visual acuity both in Pseudophakic and phakic, 6/18 were in 5 (6%) pseudophakic eyes, 6/12 in 8 (10%), 6/9 in 32 (40%) and 6/6 in 3 (44%) eyes. But in phakic 6/18 were 3 (4%), 6/12 in 5 (6%), 6/9 in 14 (18%) and 6/6 in 58 (73%) eyes as shown in Table 3.

Table 3: Distance Visual acuity

Visual acuity	Groups				Total Eyes
	pseudophakic	Percentage %	Phakic	Percentage %	
6/18	5	(6%)	3	(4%)	8
6/12	8	(10%)	5	(6%)	13
6/9	32	(40%)	14	(18%)	46
6/6	3	(44%)	58	(73%)	93
Total	80	(100%)	80	(100%)	160

The near vision in both groups pseudophakic and phakic; N6 in 45 (52%) pseudophakic eyes and N8 in 35(47%) pseudophakic eyes, but in phakic 41 (48%) eyes with N6 and 39 (53%) eyes with N8 as shown in Table 4.

Table 4: Near Visual Acuity

Group	Near Vision				Total Eyes
	N6	Percentage %	N8	Percentage %	
pseudophakic	45	52%	35	47 %	80
Phakic	41	48 %	39	53 %	80
Total	86	100 %	74	100 %	160

In Pseudophakic group myopes were 15 (19%), hypermetropes 12 (15%), astigmatic 32 (40%) and 21 (26%) had no refractive error. In phakic group myopes were 12 (15%), hypermetropes 7 (9%), astigmatic 17 (21%) and 44 (55%) had no refractive error as shown in Figure 1.




Fig 1: Type of refractive errors in Pseudophakic and phakic groups

Among pseudophakic group; tritanopia in 32 (40%), deuteronopia in 10 (13%), protonopia in 8 (10%), combined tritanopia + protonopia in 6 (8%), combined deutran + tritan in 4 (5%), combined protan + deutran in 2 (3%) eyes while 18 (23%) eyes showed no color vision defect as shown in Figure 2.




Fig 2: Defect of color vision in Pseudophakic Group

In phakic group: Tritanopia in 7 (9%) eyes and Deuteronopia 1(1.00%) while 72 (90%) had no defect as shown in Figure 3.




Fig 3: Defect of color vision in phakic Group

The below Figure 4 shows comparison between pseudophakic and phakic groups, the pseudophakic eyes were more sensitive to tritanopia 32 (40%) as compared to phakic 7 (9%), than deuteronopia 10 (13%) in pseudophakic group and only 1(9%) in phakic group, protonopia 8(10%) in pseudophakic and 0 (zero) in phakic and the combined tritanopia+protonopia defect 6 (8%) in pseudophakic but 0 (zero) in phakic, then combined deuteronopia + tritanopia 4 (5%) in pseudophakic and 0 (zero) in phakic group, combined protonopia + deuteronopia 2 (3%) in pseudophakic and 0 (zero) in phakic while 18 (23%) in pseudophakic and 72 (90%) in phakic group has no color vision defect.




Fig 4: Color vision defect in Pseudophakic Group vs Phakic Group

Discussion

In present study Pseudophakic group was more sensitive to tritanopia which contradict with the results of another study where anomaloscope and the 100-hue test were used indicating that the pseudophakic eyes were more sensitive to red and less sensitive to blue than healthy phakic eyes^[15].

In a study carried out on sixty-eight eyes of 40 diabetic patients, divided into four subgroups at different stages of diabetic retinopathy and 20 eyes of 10 healthy individuals as controls showed that the Ishihara pseudo iso-chromatic plates test; only 51% of diabetic patients passed the test, 28% failed and the remaining 21% were suspects while 90% of controls passed and only 10% failed. Only 10% of controls failed the Farnsworth D-15 test due to Protanopia, while 50% of the diabetics failed the test, with variable dyschromatopsia mainly Tritanopia and combined color vision deficiencies^[16]. Contrarily our study excluded systemic diseases.

Another cross-sectional study used the Farnsworth 100 hue test and Pickford Nicholson anomaloscope in pseudophakic, phakic and spectacle aphakic eyes to determine the little difference in their color perception. The pseudophakic eyes are highly sensitive to red and low sensitive to blue when compared with aphakic while in our study Panel D15 was used to assess color vision defect and shared contrary results showing that pseudophakic eyes were sensitive to blue.^[17]

A study showed that blue-yellow defects were becoming increasingly prevalent with increasing age^[18] similar to our study's results. Another study compared color differentiation of 30 phakic and 30 pseudophakic eyes, using the Farnsworth-Munsell 100-hue test and they found no significant difference between the two groups as regards differentiation of colors although theoretically it could be expected that color differentiation will be better in eyes with a synthetic intraocular lens and subject's age^[19]. has greatest influence on color sense while our study showed that pseudophakic group is more sensitive to blue defect as compared to phakic group.

Conclusion

The study concluded that tritanopia was more commonly present in Pseudophakic group while majority of subjects in Phakic group did not show color vision defect.

Acknowledgement

Special thanks to

Professor Dr. Mohammad Saleh Memon

Conflict of interest

None

References

- Dain S. Clinical color vision tests. *Clin. Exp. Optom.* 2004; 87:276-293.
- Roy MS, Podgor MJ, Collier B, Gunkel RD. Color vision and age in a normal North American population. *Graefe's Arch Clin Exp Ophthalmol.* 1991; 229:139-144.
- Drance SM, Lakowski R, Schulzer M, Douglas GR. Acquired color vision changes in glaucoma. *Arch Ophthalmol.* 1981; 99:829-831.
- Sehlapelo R, Oduntan A. Effect of optical defocus on colour perception. *S Afr Optom.* 2007; 66(2):77-81.
- Paramei G, Bimler D. Colour vision testing. In: *Encyclopedia of color science and technology.* New York: Springer Science and Business Media. 2015: 1-7.
- Birch J. Worldwide prevalence of red-green color deficiency. *J Opt Soc Am A Opt Image Sci Vis.* 2012; 29(3):313-320.
- Gundogan NU, Durmazlar N, Gümüş K. Projected color slides as a method for mass screening test for color vision deficiency (a preliminary study). *Int J Neurosci.* 2005; 115(8):1105-1117.
- Gordon N. Colour deficiency. *Public Health.* 1998; 112(2):81-84
- Diane VS, Fatima NM, Sershani G, Lunga L, Benita S, Osman A, *et al.* Comparing the validity of an online Ishihara colour vision test to the traditional Ishihara handbook in a South African university population. *African vision and eye health.* 2018; 77(1):1-4.
- Birch J. Worldwide prevalence of red-green color deficiency. *J Opt Soc Am A Opt Image Sci Vis.* 2012; 29(3):313-20.
- Rogosic V, Bojic L, Karaman K, Rogosic LV, Titlic M, Poljak NK, *et al.* Comparative follow-up study of unselected male population with congenital defective color vision from inland and Mediterranean areas of Croatia. *Acta Med Croatica.* 2011; 65(1):19-24.
- Shaukat AC, Farzeen KH, Shehreen A, Mustafa K, Khabir A. Frequency of color blindness in pre-employment screening in a tertiary health care center in Pakistan. *Pak J med Sci.* 2017; 33(2):430-432
- Hamida Tehmina S, Amna B, Nabeela T, Naheed S, Kashif U, *et al.* Incidence of Protanopia and Deuteranopia, Defects of Colour Vision in Quetta, Pakistan. *Pakistan J. Zool.* 2016; 48(4):1045-1049.
- Siddiqui QA, Shaikh SA, Qureshi TZ, Subhan MM. A comparison of red-green color vision deficiency between medical and non-medical students in Pakistan. *Saudi Med J.* 2010; 31(8):895899.
- Jeffrey LJ, Vijay BG, Donald A. Colour perception in pseudophakia. *British Journal of Ophthalmology.* 1982; 66:658-662.
- Tarek MR, Ehab MG, Waleed AG, Ahmed AO. Assessment of Color vision in diabetic patients. *International Journal of ophthalmic research.* 2015; 1(1):19-23.
- Jay JL, Guatam VB, Allan D. Color perception in pseudophakia. *British Journal of Ophthalmology.* 1982; 66:658-62.

18. Hassan H, Mehdi K, Reza P, Abbasali Y. The prevalence of color vision deficiency in the northeast of Iran. *Journal of current Ophthalmology*, 2017, 80-85.
19. Marilyn ES, Gunilla HP, Lori AL, John AB. Comparison of Panel D-15 Tests in a Large Older Population. *Optom Vis sci*. 2014; 91(3):284-90.
20. Skorkovska S, Skorkovsky, Janula J. Color Vision in Pseudophakia. *Cesk Oftalmol*. 1992; 48(1):10-14.